


Exhibit 8

Abstract

Degrave *et al.*, Mol Biol Rep, 11:57-61 (1986)



PubMed Nucleotide Protein Genome Structure PMC Taxonomy OMIM Bc
Search PubMed for
Limits Preview/Index History Clipboard Details

About Entrez

Show: 20

Text Version

☐ 1: Mol Biol Rep 1986;11(1):57-61

Links

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Browser

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

Privacy Policy

Cloning and structure of a mouse interleukin-2 chromosomal gene.

Degrave W, Simons G, Devos R, Plaetinck G, Remaut E, Tavernier J, Fiers W.

Using non-stringent hybridization with a human interleukin-2 cDNA probe, we have isolated recombinant phages from a mouse genomic DNA library cloned in the EMBL3 phage. The sequence and organization of the mouse interleukin-2 (IL-2) gene was determined. By comparison with the human IL-2 sequence, three introns can be identified with lengths of 99, +/- 2 400, and +/- 1 900 base pairs, respectively. The mouse IL-2 gene codes for a polypeptide of 169 amino acids and contains a putative signal peptide of 20 amino acids. The homology to the human interleukin-2 is 72% at the nucleotide level in the coding part and 65% at the amino acid level. An extraordinary sequence, consisting of 12 consecutive CAG codons coding for glutamine, is found in the first exon.

PMID: 3003564 [PubMed - indexed for MEDLINE]

Show: 20

[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Freedom of Information Act](#) | [Disclaimer](#)

i686-pc-linux-gnu Jan 7 2003 16:40:32